

Seiberg-Witten Flow in Higher Dimensions

Lorenz Schabrun *

Abstract. We show that for manifolds of dimension $m \geq 5$, the flow of a Seiberg-Witten-type functional admits a global smooth solution.

1 Introduction

The Seiberg-Witten invariant has proven a very effective tool in four-dimensional geometry. Its computation involves finding non-trivial solutions to the system of first order Seiberg-Witten equations, called monopoles. Monopoles represent the zeros of the Seiberg-Witten functional (2) (see [9]). In [6], the flow for the Seiberg-Witten functional on a 4-manifold was studied. It was shown that the flow admits a global solution which converges in C^∞ to a critical point of the functional.

The Seiberg-Witten equations and functional do not generalize immediately to higher dimensions, since they depend on the notion of self-duality on four dimensional manifolds. Nonetheless, a number of generalizations of Seiberg-Witten theory have been suggested for higher dimensional manifolds, see for example [5], [1] and [3]. In this paper, we extend the global existence result obtained for the Seiberg-Witten functional in [6] for dimension 4 to a functional of similar form in higher dimensions.

The main additional estimate in establishing global existence in higher dimensions is a so-called monotonicity formula. This idea was used by Struwe for the heat flow of harmonic maps in higher dimensions [13], and has also been used to study the Yang-Mills and Yang-Mills-Higgs flows in higher dimensions, see [2] and [7]. See also [10] and [11] for the harmonic map flow, and [14] for sequences of weakly converging Yang-Mills connections.

Let M be a compact oriented Riemannian m -manifold which admits a Spin^c structure $\mathfrak{s}$. Denote by $\mathcal{S} = W \otimes \mathcal{L}$ the corresponding spinor bundle and by $\mathcal{S}^\pm = W^\pm \otimes \mathcal{L}$ the half spinor bundles, and by $\mathcal{L}^2$ the corresponding determinant line bundle. Recall that the bundle $\mathcal{S}^+$ has fibre $\mathbb{C}^2$. Let A be a unitary connection on $\mathcal{L}^2$. Note that we can write $A = A_0 + a$, where A_0 is some fixed connection and $a \in i\Lambda^1 M$ with $i = \sqrt{-1}$. Denote by $F_A = dA \in i\Lambda^2 M$ the curvature of the line bundle connection A . Let $\{e_j\}$ be an orthonormal basis of $\mathbb{R}^4$. A

*Department of Mathematics, The University of Queensland, Brisbane, QLD 4072, Australia. Email: lorenz@maths.uq.edu.au

$\text{Spin}(4)^c$ -connection on the bundles $\mathcal{S}$ and $\mathcal{S}^\pm$ is locally defined by

$$\nabla_A = d + \frac{1}{2}(\omega + A), \quad (1)$$

where $\omega = \frac{1}{2}\omega_{jk}e_j e_k$ is induced by the Levi-Civita connection matrix ω_{jk} and $e_j e_k$ acts by Clifford multiplication (see [8]). We denote the curvature of ∇_A by Ω_A . We define the configuration space $\Gamma(\mathcal{S}^+) \times \mathcal{A}$, where $\mathcal{A}$ is the space of unitary connections on $\mathcal{L}^2$, and let $(\varphi, A) \in \Gamma(\mathcal{S}^+) \times \mathcal{A}$.

We first recall the definition of the Seiberg-Witten functional on 4-manifolds. The Seiberg-Witten functional $\mathcal{SW} : \Gamma(\mathcal{S}^+) \times \mathcal{A} \rightarrow \mathbb{R}$ is given by

$$\mathcal{SW}(\varphi, A) = \int_M |\nabla_A \varphi|^2 + |F_A^+|^2 + \frac{S}{4}|\varphi|^2 + \frac{1}{8}|\varphi|^4 dV, \quad (2)$$

where S is the scalar curvature of M . The Seiberg-Witten functional (2) is invariant under the action of a gauge group. The group of gauge transformations is

$$\mathcal{G} = \{g : M \rightarrow U(1)\}.$$

$\mathcal{G}$ acts on elements of the configuration space via

$$g^*(\varphi, A) = (g^{-1}\varphi, A + 2g^{-1}dg).$$

Using the relation

$$\|F_A\|_{L^2}^2 = 2\|F_A^+\|_{L^2}^2 - 4\pi^2 c_1(\mathcal{L})^2, \quad (3)$$

where $c_1(\mathcal{L})$ is the first Chern class of $\mathcal{L}$ (see [12]), one can also write the functional in the form

$$\mathcal{SW}(\varphi, A) = \int_M |\nabla_A \varphi|^2 + \frac{1}{2}|F_A|^2 + \frac{S}{4}|\varphi|^2 + \frac{1}{8}|\varphi|^4 dV + \pi^2 c_1(\mathcal{L})^2. \quad (4)$$

Now, consider again the case of an m -manifold M . The functional (2) is not defined here, since self-duality is a phenomenon that occurs only in dimension 4. However, we may use (4) to extend the Seiberg-Witten functional to higher dimensions. Note that the constant term $\pi^2 c_1(\mathcal{L})^2$ does not affect the Euler-Lagrange equations and so is irrelevant for the results in this paper. The Euler-Lagrange equations for the Seiberg-Witten functional are

$$-\nabla_A^* \nabla_A \varphi - \frac{1}{4}[S + |\varphi|^2]\varphi = 0, \quad (5)$$

$$-d^* F_A - i \operatorname{Im} \langle \nabla_A \varphi, \varphi \rangle = 0, \quad (6)$$

and as in [6], we define the flow of the Seiberg-Witten functional by

$$\frac{\partial \varphi}{\partial t} = -\nabla_A^* \nabla_A \varphi - \frac{1}{4}[S + |\varphi|^2]\varphi, \quad (7)$$

$$\frac{\partial A}{\partial t} = -d^* F_A - i \operatorname{Im} \langle \nabla_A \varphi, \varphi \rangle \quad (8)$$

with initial data

$$(\varphi(0), A(0)) = (\varphi_0, A_0).$$

Regarding the existence of solutions to the flow (7) and (8), we prove the following theorem.

Theorem 1. *For any given smooth (φ_0, A_0) and m -dimensional Riemannian manifold M for $m \geq 5$, equations (7) and (8) admit a unique global smooth solution on $M \times [0, \infty)$ with initial data (φ_0, A_0) .*

As in [6], we have the energy inequality

$$\frac{d}{dt} \mathcal{SW}(\varphi(t), A(t)) = - \int_M \left[2 \left| \frac{\partial \varphi}{\partial t} \right|^2 + \left| \frac{\partial A}{\partial t} \right|^2 \right] \leq 0. \quad (9)$$

or

$$\int_0^T \left[2 \left\| \frac{\partial \varphi}{\partial t} \right\|_{L^2}^2 + \left\| \frac{\partial A}{\partial t} \right\|_{L^2}^2 \right] = \mathcal{SW}(\varphi_0, A_0) - \mathcal{SW}(\varphi(T), A(T)). \quad (10)$$

Many of the proofs in [6] do not contain dimensional considerations, and are also valid in the m -dimensional case. For the proofs of the following Lemmas, we direct the reader to that paper.

Lemma 1.1. *For any given smooth initial data (φ_0, A_0) , the system (7)-(8) admits a unique local smooth solution on $M \times [0, T)$ for some $T > 0$.*

Lemma 1.2. *Let (φ, A) be a solution of (7)-(8) on $M \times [0, T)$, and write $m = \sup_{x \in M} |\varphi_0|$. Then for all $t \in [0, T)$, we have*

$$\sup_{x \in M} |\varphi(x, t)| \leq \max\{m, \sqrt{|S_0|}\}. \quad (11)$$

Lemma 1.3. *There exist positive constants c, c' such that the following estimate holds:*

$$\begin{aligned} & \frac{\partial}{\partial t} (|\nabla_A \varphi|^2 + |F_A|^2) + \Delta (|\nabla_A \varphi|^2 + |F_A|^2) \\ & \leq -c' (|\nabla_A^2 \varphi|^2 + |\nabla F_A|^2) + c(|F_A| + 1) (|\nabla_A \varphi|^2 + |F_A|^2 + 1). \end{aligned}$$

Lemma 1.4. *Let (φ, A) be a solution to (7)-(8) in $M \times [0, T)$ with initial values (φ_0, A_0) . Suppose $|\nabla_A \varphi| \leq K_1$ and $|F_A| \leq K_1$ in $M \times [0, T)$ for some constant $K_1 > 0$. Then for any positive integer $n \geq 1$, there is a constant K_{n+1} independent of T such that*

$$\left| \nabla_A^{(n+1)} \varphi \right| \leq K_{n+1}, \quad \left| \nabla_M^{(n)} F_A \right| \leq K_{n+1} \quad \text{in } M \times [0, T),$$

where (n) denotes n iterations of the derivative.

2 Estimates

We first derive a monotonicity inequality for the flow (7)-(8). We define

$$e(\varphi, A)(x, t) = |\nabla_A \varphi|^2 + \frac{1}{2} |F_A|^2 + \frac{S}{4} |\varphi|^2 + \frac{1}{8} |\varphi|^4.$$

Let $z = (x, t)$ denote a point of $M \times \mathbb{R}$, with $z_0 = (x_0, t_0) \in M \times [0, T]$. We define

$$T_R(z_0) = M \times [t_0 - 4R^2, t_0 - R^2],$$

and

$$P_R(z_0) = B_R(x_0) \times [t_0 - R^2, t_0],$$

where $B_R(x_0) \subset M$ denotes a ball centered at x_0 with radius R . Note that in constructing $T_R(z_0)$ we require that $t_0 - 4R^2 \geq 0$ or $R \leq \sqrt{t_0}/2$. We abbreviate $T_R(0, 0) = T_R$ and $P_R(0, 0) = P_R$. The fundamental solution to the backward heat equation with singularity at z_0 is

$$G_{z_0}(z) = \frac{1}{(4\pi(t_0 - t))^{m/2}} \exp\left(-\frac{(x - x_0)^2}{4(t_0 - t)}\right)$$

where $t < t_0$. We also write $G = G_{(0,0)}$. Let $i(M)$ be the injectivity radius of M , and suppose that (φ, A) is a solution to the flow (7)-(8) on $M \times [0, T]$. Let ϕ_x be a smooth cut-off function with $|\phi_x| \leq 1$, $\phi_x \equiv 1$ on $B_{i(M)/2}(x)$, $\phi_x \equiv 0$ outside $B_{i(M)}(x)$ and $|\nabla \phi_x| \leq c/i(M)$ for some constant c . We also abbreviate $\phi = \phi_{x_0}$. We define

$$\Phi(R; \varphi, A) = R^2 \int_{T_R(z_0)} e(\varphi, A)(z) \phi^2 G dV dt, \quad (12)$$

and

$$\mathcal{F}(R; \varphi, A) = \int_{T_R(z_0)} Rt \left(\left| \frac{\partial A}{\partial t} + \frac{x_k}{2t} \frac{\partial}{\partial x_k} \right| F_A \right|^2 + 2 \left| \frac{\partial \varphi}{\partial t} + \frac{x_k}{2t} \nabla_A^k \varphi \right|^2 \right) \phi^2 G \sqrt{g} dz, \quad (13)$$

where

$$\frac{\partial}{\partial x_k} \rfloor F_A = F_A \left(\frac{\partial}{\partial x_k}, \cdot \right) = F^{kj} dx^j$$

defines a 1-form.

Lemma 2.1. *Let (φ, A) be a smooth solution of (7)-(8) on $M \times [0, T]$ with initial data (φ_0, A_0) . Then for $z_0 \in M \times [0, T]$ and any R_a and R_b satisfying $0 < R_a \leq R_b \leq R_0$ for some $R_0 \leq \min\{i(M), \sqrt{t_0}/2\}$, we have*

$$\Phi(R_a; \varphi, A) + \int_{R_a}^{R_b} e^{aR} F(R) \leq e^{c(R_b - R_a)} \Phi(R_b; \varphi, A) + c(R_b^2 - R_a^2) \mathcal{SW}(\varphi_0, A_0),$$

where c depends only on the geometry of M .

Proof. We show that

$$\frac{d}{dR}\Phi(R; \varphi, A) + \mathcal{F}(R; \varphi, A) \geq -c\Phi(R; \varphi, A) - cRSW(\varphi_0, A_0). \quad (14)$$

The required result then follows by multiplying (14) by e^{aR} for some sufficiently large $a > 0$, and integrating from R_a to R_b . To show (14), we may assume that $z_0 = (0, 0)$, which implies that $t < 0$ on T_R . We rescale the coordinates to $x = R\tilde{x}$, $t = R^2\tilde{t}$. In these coordinates,

$$\Phi(R; \varphi, A) = \int_{T_1} R^4 e(\varphi, A)(R\tilde{x}, R^2\tilde{t}) \phi^2(R\tilde{x}) G(\tilde{z}) \sqrt{g(R\tilde{x})} d\tilde{z}$$

where $d\tilde{z} = d\tilde{x}d\tilde{t}$. For some $R \leq R_0$, we compute

$$\begin{aligned} \frac{d}{dR}\Phi(R; \varphi, A) &= \int_{T_1} \frac{d}{dR} \left[R^4 e(\varphi, A)(R\tilde{x}, R^2\tilde{t}) \phi^2(R\tilde{x}) \sqrt{g(R\tilde{x})} \right] G(\tilde{z}) d\tilde{z} \\ &= \int_{T_1} 4R^3 e(\varphi, A)(R\tilde{x}, R^2\tilde{t}) \phi^2(R\tilde{x}) G(\tilde{z}) \sqrt{g(R\tilde{x})} d\tilde{z} \\ &\quad + \int_{T_1} R^4 \tilde{x}_k \frac{\partial}{\partial x_k} e(\varphi, A)(R\tilde{x}, R^2\tilde{t}) \phi^2(R\tilde{x}) G(\tilde{z}) \sqrt{g(R\tilde{x})} d\tilde{z} \\ &\quad + \int_{T_1} 2R^5 \tilde{t} \frac{\partial}{\partial t} e(\varphi, A)(R\tilde{x}, R^2\tilde{t}) \phi^2(R\tilde{x}) G(\tilde{z}) \sqrt{g(R\tilde{x})} d\tilde{z} \\ &\quad + \int_{T_1} R^4 e(\varphi, A)(R\tilde{x}, R^2\tilde{t}) \tilde{x}_k \frac{\partial}{\partial x_k} (\phi^2 \sqrt{g})(R\tilde{x}) G(\tilde{z}) d\tilde{z} \\ &:= I_1 + I_2 + I_3 + I_4. \end{aligned}$$

Rescaling coordinates back to (x, t) , we have

$$I_1 = \int_{T_R} 4Re(\varphi, A) \phi^2 G \sqrt{g} dz$$

and

$$I_4 = \int_{T_R} Re(\varphi, A) x_k \frac{\partial}{\partial x_k} (\phi^2 \sqrt{g}) G dz.$$

For the second term,

$$I_2 = \int_{T_R} R x_k \frac{\partial}{\partial x_k} e(\varphi, A) \phi^2 G \sqrt{g} dz.$$

This simplifies as follows:

$$\begin{aligned} &\frac{\partial}{\partial x_k} \left[|\nabla_A \varphi|^2 + \frac{1}{2} |F_A|^2 + \left(\frac{S}{4} |\varphi|^2 + \frac{1}{8} |\varphi|^4 \right) \right] \\ &= \langle \nabla_M^k F_A, F_A \rangle + 2 \operatorname{Re} \langle \nabla_A^k \nabla_A^j \varphi, \nabla_A^j \varphi \rangle + \frac{1}{2} (S + |\varphi|^2) \operatorname{Re} \langle \nabla_A^k \varphi, \varphi \rangle. \end{aligned}$$

Note that

$$2 \operatorname{Re} \langle \nabla_A^k \nabla_A^j \varphi, \nabla_A^j \varphi \rangle = 2 \operatorname{Re} \langle \nabla_A \nabla_A^k \varphi, \nabla_A \varphi \rangle - 2 \operatorname{Re} \langle \Omega_A^{jk} \varphi, \nabla_A^j \varphi \rangle.$$

Using the fact that

$$\frac{\partial G}{\partial x_j} = \frac{x_j}{2t} G,$$

we have

$$-2 \int_{T_R} R x_k \operatorname{Re} \langle d(G) \wedge \nabla_A^k \varphi, \nabla_A \varphi \rangle \phi^2 \sqrt{g} dz = -4 \int_{T_R} R t \left| \frac{x_k}{2t} \nabla_A^k \varphi \right|^2 \phi^2 G \sqrt{g} dz.$$

For the curvature term, we recall the Bianchi identity $dF_A = 0$ which implies that

$$\partial_k F^{ij} = \partial_i F^{kj} - \partial_j F^{ki},$$

and we compute in local coordinates

$$\begin{aligned} x_k \partial_k \sum_{i < j} (F^{ij})^2 G \phi^2 &= 2x_k \sum_{i < j} F^{ij} \partial_k F^{ij} G \phi^2 \\ &= 2x_k \sum_{i < j} F^{ij} (\partial_i F^{kj} - \partial_j F^{ki}) G \phi^2 \\ &= x_k F^{ij} \partial_i F^{kj} G \phi^2 \\ &= \partial_i (x_k F^{ij} F^{kj} G \phi^2) - x_k F^{kj} \partial_i F^{ij} G \phi^2 - F^{kj} F^{ij} \partial_i (x_k G \phi^2). \end{aligned}$$

Observe that the first term will integrate to zero by Stoke's theorem, and for the second term we have $(d^* F_A)_j = \partial_i F^{ij}$. To deal with the third term, we see that

$$-x_k F^{kj} x_i F^{ij} \frac{1}{2t} G \phi^2 = -2t \left| \frac{x_k}{2t} \frac{\partial}{\partial x_k} F_A \right|^2 G \phi^2.$$

Note that $|d\phi| G \leq c$ since $|d\phi| = 0$ on $B_{i(M)/2}(x_0)$. Then

$$\begin{aligned} I_2 &= \int_{T_R} R x_k \left[\frac{1}{2} \partial_k |F_A|^2 + 2 \operatorname{Re} \langle \nabla_A \nabla_A^k \varphi, \nabla_A \varphi \rangle - 2 \operatorname{Re} \langle \Omega_A^{jk} \varphi, \nabla_A^j \varphi \rangle \right. \\ &\quad \left. + \frac{1}{2} (S + |\varphi|^2) \operatorname{Re} \langle \nabla_A^k \varphi, \varphi \rangle \right] \phi^2 G \sqrt{g} dz \\ &= \int_{T_R} R x_k \left[\frac{1}{2} \partial_k |F_A|^2 + 2 \operatorname{Re} \langle \nabla_A^k \varphi, \nabla_A^* \nabla_A \varphi \rangle \right. \\ &\quad \left. - 2 \operatorname{Re} \langle \Omega_A^{jk} \varphi, \nabla_A^j \varphi \rangle + \frac{1}{2} (S + |\varphi|^2) \operatorname{Re} \langle \nabla_A^k \varphi, \varphi \rangle \right] \phi^2 G \sqrt{g} dz \\ &= 2 \int_{T_R} R x_k \operatorname{Re} \langle d(\phi^2 G x_k) \nabla_A^k \varphi, \nabla_A \varphi \rangle \sqrt{g} dz \\ &\geq - \int_{T_R} R |x| \left(\left| \frac{\partial A}{\partial t} \right| |F_A| + 2 \left| \frac{\partial \varphi}{\partial t} \right| |\nabla_A \varphi| \right) \phi^2 G \sqrt{g} dz \\ &\quad - 4 \int_{T_R} R t \left| \frac{x_k}{2t} \nabla_A^k \varphi \right|^2 \phi^2 G \sqrt{g} dz - \int_{T_R} R t \left| \frac{x_k}{2t} \frac{\partial}{\partial x_k} F_A \right|^2 G \phi^2 \sqrt{g} dz \\ &\quad - c \Phi(R; \varphi, A) - c R \mathcal{SW}(\varphi_0, A_0), \end{aligned}$$

where we note that $\Omega_A = \Omega_{A_0} + \frac{1}{2} F_A$, and we also recall the fact from [6] that

$$\operatorname{Re} \left\langle \frac{\partial A}{\partial t} \varphi, \nabla_A \varphi \right\rangle = \left\langle \frac{\partial A}{\partial t}, i \operatorname{Im} \langle \nabla_A \varphi, \varphi \rangle \right\rangle;$$

note that $\frac{\partial A}{\partial t}$ can be replaced by any 1-form. For the third term,

$$\begin{aligned}
I_3 &= \int_{T_R} 2Rt \left[\left\langle d \frac{\partial A}{\partial t}, F_A \right\rangle + 2 \operatorname{Re} \left\langle \nabla_A \frac{\partial \varphi}{\partial t}, \nabla_A \varphi \right\rangle + \operatorname{Re} \left\langle \frac{\partial A}{\partial t} \varphi, \nabla_A \varphi \right\rangle \right. \\
&\quad \left. + \left(\frac{1}{2} [S + |\varphi|^2] \operatorname{Re} \left\langle \frac{\partial \varphi}{\partial t}, \varphi \right\rangle \right) \right] \phi^2 G \sqrt{g} dz \\
&= - \int_{T_R} 2Rt \left[\left| \frac{\partial A}{\partial t} \right|^2 + 2 \left| \frac{\partial \varphi}{\partial t} \right|^2 \right] \phi^2 G \sqrt{g} dz \\
&\quad - \int_{T_R} 2Rt \left\langle d(\phi^2 G) \wedge \frac{\partial A}{\partial t}, F_A \right\rangle \sqrt{g} dz \\
&\quad - \int_{T_R} 4Rt \operatorname{Re} \left\langle d(\phi^2 G) \wedge \frac{\partial \varphi}{\partial t}, \nabla_A \varphi \right\rangle \sqrt{g} dz.
\end{aligned}$$

Next we obtain

$$\left\langle dG \wedge \frac{\partial A}{\partial t}, F_A \right\rangle = \left\langle \frac{x_k}{2t} dx^k \wedge \frac{\partial A}{\partial t}, F_A \right\rangle G = \left\langle \frac{\partial A}{\partial t}, \frac{x_k}{2t} \frac{\partial}{\partial x_k} \lrcorner F_A \right\rangle G,$$

and

$$\left\langle dG \wedge \frac{\partial \varphi}{\partial t}, \nabla_A \varphi \right\rangle = \left\langle \frac{x_k}{2t} dx^k \wedge \frac{\partial \varphi}{\partial t}, \nabla_A \varphi \right\rangle G = \left\langle \frac{\partial \varphi}{\partial t}, \frac{x_k}{2t} \nabla_A^k \varphi \right\rangle G.$$

Thus

$$\begin{aligned}
I_3 &\geq - \int_{T_R} 2Rt \left[\left| \frac{\partial A}{\partial t} \right|^2 + 2 \left| \frac{\partial \varphi}{\partial t} \right|^2 \right] \phi^2 G \sqrt{g} dz \\
&\quad - \int_{T_R} 2Rt \left\langle \frac{\partial A}{\partial t}, \frac{x_k}{2t} \frac{\partial}{\partial x_k} \lrcorner F_A \right\rangle \phi^2 G \sqrt{g} dz + c \int_{T_R} 2Rt \left| \frac{\partial A}{\partial t} \right| |F_A| \phi \sqrt{g} dz \\
&\quad - \int_{T_R} 4Rt \operatorname{Re} \left\langle \frac{\partial \varphi}{\partial t}, \frac{x_k}{2t} \nabla_A^k \varphi \right\rangle \phi^2 G \sqrt{g} dz + c \int_{T_R} 4Rt \left| \frac{\partial \varphi}{\partial t} \right| |\nabla_A \varphi| \phi \sqrt{g} dz. \\
&\geq - \int_{T_R} Rt \left(\left| \frac{\partial A}{\partial t} + \frac{x_k}{2t} \frac{\partial}{\partial x_k} \lrcorner F_A \right|^2 + 2 \left| \frac{\partial \varphi}{\partial t} + \frac{x_k}{2t} \nabla_A^k \varphi \right|^2 \right) \phi^2 G \sqrt{g} dz \\
&\quad + \int_{T_R} Rt \left(\left| \frac{x_k}{2t} \frac{\partial}{\partial x_k} \lrcorner F_A \right|^2 + 2 \left| \frac{x_k}{2t} \nabla_A^k \varphi \right|^2 \right) \phi^2 G \sqrt{g} dz \\
&\quad - \int_{T_R} Rt \left[\left| \frac{\partial A}{\partial t} \right|^2 + 2 \left| \frac{\partial \varphi}{\partial t} \right|^2 \right] \phi^2 G \sqrt{g} dz \\
&\quad - cRS\mathcal{W}(\varphi_0, A_0)
\end{aligned}$$

using Young's inequality, the energy inequality (10), and noting that $|t| \leq 4R^2$ and $R \leq R_0$. Finally, since as in [2], $R^{-1} |x|^2 G \leq c(1+G)$, combining the working above (and recalling that $t < 0$ on T_R), one obtains (14). $\square$

Corollary 2.2. *There exists a constant $a > 0$ such that*

$$e^{aR} \Phi(R; \varphi, A) + cR^2 S\mathcal{W}(\varphi_0, A_0)$$

is non-decreasing in R , where c here represents the same constant which appears in (14).

Proof. The result follows from (14) by multiplying by e^{aR} for some sufficiently large $a > 0$, and integrating from R_a to R_b . $\square$

Lemma 2.3. *Suppose $(\varphi, A) \in C^\infty(P_R(y, s))$ satisfies (7)-(8). Then there exist constants δ and R_1 such that if $R \leq R_1$ and*

$$\sup_{0 < t < s} R^{4-m} \int_{B_R(y)} (|\nabla_A \varphi|^2 + |F_A|^2) dV < \delta,$$

then

$$\sup_{P_{R/2}(y, s)} (|\nabla_A \varphi|^2 + |F_A|^2) \leq 256R^{-4}.$$

Proof. We begin by choosing $r_0 < R$ so that

$$(R - r_0)^4 \sup_{P_{r_0}(y, s)} (|\nabla_A \varphi|^2 + |F_A|^2) = \max_{0 \leq r \leq R} \left[(R - r)^4 \sup_{P_r(y, s)} (|\nabla_A \varphi|^2 + |F_A|^2) \right]. \quad (15)$$

Let

$$e_0 = \sup_{P_{r_0}(y, s)} (|\nabla_A \varphi|^2 + |F_A|^2) = (|\nabla_A \varphi|^2 + |F_A|^2)(x_0, t_0)$$

for some $(x_0, t_0) \in \bar{P}_{r_0}(y, s)$. We claim that

$$e_0 \leq 16(R - r_0)^{-4}. \quad (16)$$

Then

$$\begin{aligned} (R - r)^4 \sup_{P_r(y, s)} (|\nabla_A \varphi|^2 + |F_A|^2) &\leq (R - r_0)^4 \sup_{P_{r_0}(y, s)} (|\nabla_A \varphi|^2 + |F_A|^2) \\ &\leq 16(R - r_0)^4 (R - r_0)^{-4} = 16 \end{aligned}$$

for any $r < R$. Choosing $r = \frac{1}{2}R$ in the above, we have the required result.

We now prove (16). Define $\rho_0 = e_0^{-1/4}$ and suppose by contradiction that $\rho_0 \leq \frac{1}{2}(R - r_0)$. We rescale variables $x = x_0 + \rho_0 \tilde{x}$ and $t = t_0 + \rho_0^2 \tilde{t}$ and set

$$\psi(\tilde{x}, \tilde{t}) = \varphi(x_0 + \rho_0 \tilde{x}, t_0 + \rho_0^2 \tilde{t}),$$

$$B(\tilde{x}, \tilde{t}) = \rho_0 A(x_0 + \rho_0 \tilde{x}, t_0 + \rho_0^2 \tilde{t}),$$

giving

$$\begin{aligned} |\nabla_B \psi|^2 &= \rho_0^2 |\nabla_A \varphi|^2, \\ |F_B|^2 &= \rho_0^4 |F_A|^2. \end{aligned}$$

We define

$$e_{\rho_0}(\tilde{x}, \tilde{t}) = |F_B|^2 + \rho_0^2 |\nabla_B \psi|^2 = \rho_0^4 (|\nabla_A \varphi|^2 + |F_A|^2)$$

so that

$$e_{\rho_0}(\tilde{x}, \tilde{t}) \leq e_{\rho_0}(0, 0) = 1.$$

We compute

$$\begin{aligned} \sup_{\tilde{P}_1(0,0)} e_{\rho_0}(\tilde{x}, \tilde{t}) &= \rho_0^4 \sup_{P_{\rho_0}(x_0, t_0)} \left(|\nabla_A \varphi|^2 + |F_A|^2 \right) \\ &\leq \rho_0^4 \sup_{P_{\frac{R+r_0}{2}}(y, s)} \left(|\nabla_A \varphi|^2 + |F_A|^2 \right) \\ &= \rho_0^4 \left(\frac{R-r_0}{2} \right)^{-4} \left(R - \frac{R+r_0}{2} \right)^4 \sup_{P_{\frac{R+r_0}{2}}(y, s)} \left(|\nabla_A \varphi|^2 + |F_A|^2 \right) \\ &\leq \rho_0^4 \left(\frac{R-r_0}{2} \right)^{-4} (R-r_0)^4 e_0 = 16, \end{aligned}$$

where we have used that $P_{\rho_0}(x_0, t_0) \subset P_{\frac{R+r_0}{2}}(y, s)$, and to get to the last line we have used (15). This implies that

$$e_{\rho_0} = \rho_0^4 \left(|\nabla_A \varphi|^2 + |F_A|^2 \right) \leq 16$$

on $\tilde{P}_1(0, 0)$. By Lemma 1.3,

$$\left(\frac{\partial}{\partial \tilde{t}} + \Delta \right) \left(|\nabla_A \varphi|^2 + |F_A|^2 + 1 \right) \leq c(|F_A| + 1) \left(|\nabla_A \varphi|^2 + |F_A|^2 + 1 \right).$$

Then

$$\begin{aligned} \left(\frac{\partial}{\partial \tilde{t}} + \tilde{\Delta} \right) (e_{\rho_0} + \rho_0^4) &= \rho_0^6 \left(\frac{\partial}{\partial \tilde{t}} + \Delta \right) \left(|\nabla_A \varphi|^2 + |F_A|^2 \right) \\ &\leq c \rho_0^6 (|F_A| + 1) \left(|\nabla_A \varphi|^2 + |F_A|^2 + 1 \right) \end{aligned}$$

on $\tilde{P}_1(0, 0)$. Note that by assumption $\rho_0 < R$, $\rho_0^2 |F_A|$ is thus bounded by a constant. Then

$$\left(\frac{\partial}{\partial \tilde{t}} + \tilde{\Delta} \right) (e_{\rho_0} + \rho_0^4) \leq c(e_{\rho_0} + \rho_0^4)$$

for a constant $c > 0$. We apply Moser's Harnack inequality to give

$$\begin{aligned} 1 + \rho_0^4 &= e_{\rho_0}(0, 0) + \rho_0^4 \leq c \int_{\tilde{P}_1(0,0)} e_{\rho_0} d\tilde{x} d\tilde{t} + c \rho_0^4 \\ &= c \rho_0^{2-m} \int_{P_{\rho_0}(x_0, t_0)} \left(|\nabla_A \varphi|^2 + |F_A|^2 \right) dV dt + c \rho_0^4 \\ &\leq c \sup_{0 \leq t \leq s} R^{4-m} \int_{B_R(y)} \left(|\nabla_A \varphi|^2 + |F_A|^2 \right) dV + c R^4 \\ &< c \delta + c R^4, \end{aligned}$$

where we have used that $\rho_0 < R$. Now if we choose R_1 and δ sufficiently small, we have the desired contradiction. $\square$

3 Singularity analysis

Let (φ, A) be a smooth solution on $[0, T)$. Suppose there exists some $R \leq R_1$ such that

$$R^{4-m} \int_{B_R(y)} (|\nabla_A \varphi|^2 + |F_A|^2) dV < \delta,$$

for all $x_0 \in M$ and $t_0 = T$. Then by Lemma 2.3, $|\nabla_A \varphi|^2$ and $|F_A|^2$ are uniformly bounded on $M \times [0, T)$. As in [6], Using Lemma 1.4 we can show that ϕ and A are smooth at $t = T$. In conjunction with the local existence result, we can extend (φ, A) to a global smooth solution.

We define the singular set

$$\Sigma = \bigcap_{0 < R \leq R_1} \left\{ x_0 \in M : \limsup_{t \rightarrow T} R^{4-m} \int_{B_R(x_0)} (|\nabla_A \varphi|^2 + |F_A|^2) dV \geq \delta \right\}.$$

By the above discussion, $(\varphi(T), A(T))$ is smooth on $M \setminus \Sigma$. Let Σ' be defined as for Σ , but with ε_0 replaced by a smaller constant. Clearly $\Sigma \subseteq \Sigma'$. Furthermore, if $x \in M \setminus \Sigma$ then by smoothness at x , $x \in M \setminus \Sigma'$. Thus replacing ε_0 with any smaller constant defines the same set.

If $x \in M \setminus \Sigma$, then by Lemma 2.3, $B_R(x) \in M \setminus \Sigma$ for some R . Thus Σ is closed. Unlike in the 4 dimensional case [6], we cannot conclude at this point that the singular set is finite. Although it will not be needed in the proof of Theorem 1, one can instead show that Σ has finite $(m-4)$ -dimensional Hausdorff measure $\mathcal{H}^{m-4}$. Explicitly, for $x_0 \in \Sigma$,

$$\delta \leq \limsup_{t \rightarrow T} R^{4-m} \int_{B_R(x_0)} e(\varphi, A) dV \quad (17)$$

for any R . The family $\mathcal{F} = \{B_R(x_0) : x_0 \in \Sigma\}$ covers Σ , and by Vitali's covering lemma, there exists a finite subfamily $\mathcal{F}' = \{B_R(x_j)\}$ such that any two balls in $\mathcal{F}'$ are disjoint and $\{B_{5R}(x_j)\}$ covers Σ . Then using (17),

$$\begin{aligned} \sum_j (5R)^{m-4} &\leq \frac{5^m}{\delta} \sum_j \limsup_{t \rightarrow T} \int_{B_R(x_j)} (|\nabla_A \varphi|^2 + |F_A|^2) dV \\ &\leq C \mathcal{SW}(\varphi_0, A_0), \end{aligned}$$

where $\{B_{5R}(x_j)\}$ covers Σ . It follows that $\mathcal{H}^{m-4}(\Sigma)$ is finite, as claimed.

To establish Theorem 1, we show that $\Sigma = \emptyset$. Suppose by contradiction that Σ is non-empty. Since the flow is smooth on $[0, T)$, we can find sequences $x_n \in M$, $t_n \rightarrow T$, $R_n \rightarrow 0$ such that

$$\begin{aligned} \delta &> R_n^{4-m} \mathcal{SW}_{B_{R_n}(x_n)}(\varphi(t_n), A(t_n)) \\ &= \sup_{0 \leq t \leq t_n, x \in M} R_n^{4-m} \mathcal{SW}_{B_{R_n}(x)}(\varphi(t), A(t)) > \frac{\delta}{2} \end{aligned} \quad (18)$$

for each n , where $\mathcal{SW}_{B_R(x)}$ is defined by

$$\mathcal{SW}_{B_R(x)}(\varphi, A) = \int_{B_R(x)} |\nabla_A \varphi|^2 + \frac{1}{2} |F_A|^2 + \frac{S}{4} |\varphi|^2 + \frac{1}{8} |\varphi|^4.$$

By the compactness of M , passing to a subsequence we have $x_n \rightarrow x_0$ where $x_0 \in \Sigma$ by Lemma 2.3. We define the region

$$D_n = \{(y, s) : R_n y + x_n \in B_{i(M)/2}(x_n), s \in [-R_n^{-2}t_n, 0]\} =: U_n \times [-R_n^{-2}t_n, 0].$$

Note that as $n \rightarrow \infty$, $D_n \rightarrow \mathbb{R}^m \times (-\infty, 0]$. Furthermore, truncating the sequence if necessary, we can arrange that $B_{i(M)/2}(x_n) \subset B_{i(M)}(x_0)$. We rescale to

$$\varphi_n(y, s) = \varphi(R_n y + x_n, R_n^2 s + t_n),$$

$$A_n(y, s) = R_n A(R_n y + x_n, R_n^2 s + t_n).$$

which are defined on D_n . We have

$$|\nabla_{A_n} \varphi_n|^2 = R_n^2 |\nabla_A \varphi|^2,$$

$$|F_{A_n}|^2 = R_n^4 |F_A|^2.$$

If we choose our local coordinates on $B_{i(M)}(x_0)$ to be orthonormal coordinates, then the metric on the rescaled space is simply $g_{ij} = \delta_{ij}$. From (18),

$$\int_{B_1(0)} R_n^2 |\nabla_{A_n} \varphi_n|^2 + |F_{A_n}|^2 + R_n^4 \left(\frac{S}{4} |\varphi_n|^2 + \frac{1}{8} |\varphi_n|^4 \right) dy > \frac{\delta}{2} \quad (19)$$

for each n and $s = 0$. Next, from Lemma 2.3 and (18) we have

$$\sup_{D_n} \left(|\nabla_{A_n} R_n \varphi_n|^2 + |F_{A_n}|^2 \right) \leq K_1, \quad (20)$$

where K_1 is independent of n . We consider the rescaled equations

$$\frac{\partial R_m \varphi_m}{\partial s} = R_m^3 \frac{\partial \varphi}{\partial t} = -\nabla_{A_m}^* \nabla_{A_m} R_m \varphi_m - \frac{1}{4} \left[R_m^2 S + |R_m \varphi_m|^2 \right] R_m \varphi_m, \quad (21)$$

$$\frac{\partial A_m}{\partial s} = R_m^3 \frac{\partial A}{\partial t} = -d^* F_{A_m} - i \operatorname{Im} \langle \nabla_{A_m} R_m \varphi_m, R_m \varphi_m \rangle. \quad (22)$$

Noting the similarity of these equations to (7) and (8), we use (20) and results identical to Lemma (1.4) to find

$$\sup_{D_n} \left(\left| \nabla_{A_n}^{(k+1)} R_n \varphi_n \right|^2 + \left| \nabla_M^{(k)} F_{A_n} \right|^2 \right) \leq K_{k+1} \quad (23)$$

for each $k \geq 0$. Thus by a result of Uhlenbeck (theorem 1.3 in [15], see also [7]), passing to a subsequence and using an appropriate gauge we have C^∞ convergence $R_n \varphi_n \rightarrow \tilde{\varphi} = 0$ (since φ_n is bounded), $A_n \rightarrow \tilde{A}$ where $\tilde{\varphi}$ and $\tilde{A}$ are defined on $\mathbb{R}^m \times (-\infty, 0]$. Then as $n \rightarrow \infty$ in (19), we obtain

$$\int_{B_2(0)} |F_{\tilde{A}}|^2 dy \geq \frac{\delta}{2} \quad (24)$$

for $s = 0$. Since $R_n \varphi_n \rightarrow 0$, from (22) $\tilde{A}$ satisfies the equation

$$\frac{\partial \tilde{A}}{\partial s} = -d^* F_{\tilde{A}}$$

on $\mathbb{R}^m \times (-\infty, 0]$. Using the Bianchi identity $dF_{\tilde{A}} = 0$, This implies that

$$\frac{\partial F_{\tilde{A}}}{\partial s} = -\Delta F_{\tilde{A}}$$

on $\mathbb{R}^m \times (-\infty, 0]$, where $\Delta = d^*d + dd^*$. Since the solution to the heat equation converges to constant data in infinite time, the only possible solution to this equation satisfying (20) is $F_{\tilde{A}} = \text{constant}$. See for example theorem 9 of chapter 2 in [4]. In the notation of [4], choose $k = 1$ and $t = 0$, note that for us $\|u\|_{L^1(C(x,t;r))} \leq cr^{n+2}$, and let $r \rightarrow \infty$.

4 Proof of Theorem 1

As in [14] and [11], the term $\mathcal{F}(R; \varphi, A)$ in Lemma 2.1 can be used to further analyse the singularity (see for example Lemma 3.3.2 of [14]). However, we are already in a position to show that the existence of a singularity implies a contradiction. Noting that $G_{(x_n, t_n)} \geq cR_n^{-m}$ on $B_{rR_n}(x_n) \times [t_n - 4(rR_n)^2, t_n - (rR_n)^2]$, we consider for any $r \in (0, \infty)$,

$$\begin{aligned} & r^{2-m} \int_{B_r(0) \times [-4r^2, -r^2]} |F_{\tilde{A}}|^2 dy ds \\ &= \lim_{n \rightarrow \infty} (rR_n)^{2-m} \int_{B_{rR_n}(x_n) \times [t_n - 4(rR_n)^2, t_n - (rR_n)^2]} |F_A|^2 dV dt \\ &\leq c \lim_{n \rightarrow \infty} (rR_n)^2 \int_{B_{rR_n}(x_n) \times [t_n - 4(rR_n)^2, t_n - (rR_n)^2]} |F_A|^2 G_{(x_n, t_n)} dV dt \\ &\leq c \lim_{n \rightarrow \infty} (rR_n)^2 \int_{T_{rR_n}(x_0, T)} e(\varphi, A) G_{(x_0, T)} \phi^2 dV dt. \end{aligned}$$

However, the latter expression is bounded by Lemma 2.1. Thus

$$\int_{B_r(0) \times [-4r^2, -r^2]} |F_{\tilde{A}}|^2 dy ds \leq cr^{m-2}.$$

But since $|F_{\tilde{A}}|$ is constant and non-zero by (24), this implies that $r^4 \leq c$. this is impossible for r sufficiently large. This proves Theorem 1. $\square$

References

- [1] Bilge A.H., Dereli T. and Kocak S., Seiberg-Witten type monopole equations on 8-manifolds with $\text{Spin}(7)$ holonomy, as minimizers of a quadratic action, *J. High Energy Phys.* 4 (2003) 003.

- [2] Chen, Y., Shen, C.-L., Monotonicity formula and small action regularity for Yang-Mills flows in higher dimensions. *Calc. Var. PDEs* 2 (1994) 389-403.
- [3] Değirmenci, N. and Özdemir, N., Seiberg-Witten-like Equations on 7-Manifolds with G_2 -structure. *J. Nonlinear Math. Phys.* 12 (2005) no.4 457-461.
- [4] Evans, L.C, *Partial differential equations*, American Mathematical Society, 1998.
- [5] Gao, Y.-H. and Tian, G., Instantons and the monopole-like equations in eight dimensions, *J. High Energy Phys.* 5 (2000) 036.
- [6] Hong, M.-C. and Schabrun, L., Global existence for the Seiberg-Witten flow, *preprint* 2009.
- [7] Hong, M.-C. and Tian, G., Asymptotical behaviour of the Yang-Mills flow and singular Yang-Mills connections, *Math. Ann.* 330 (2004) 441-472.
- [8] Jost, J., *Riemannian Geometry and Geometric Analysis*, Springer-Verlag, 1995.
- [9] Jost, J., Peng, X. and Wang G., Variational aspects of the Seiberg-Witten functional, *Calc. Var. Partial Differential Equations* 3(1996)205-218.
- [10] Lin, F.H., Gradient estimates and blow-up analysis for stationary harmonic maps *Ann. of Math.* 149 (1999) 785-829.
- [11] Lin, F.H., and Wang, C.Y., Harmonic and quasi-harmonic spheres, *Comm. Anal. Geom.* 10 (1999) no.2 397-429.
- [12] Scorpan, A., *The Wild World of 4-Manifolds*, American Mathematical Society, 2005.
- [13] Struwe, M., On the evolution of harmonic maps in higher dimensions. *J. Diff. Geom.* 28 (1988) 485-502.
- [14] Tian, G., Gauge theory and calibrated geometry, I, *Annals of Mathematics* 151 (2000) 193-268.
- [15] Uhlenbeck, K., Connections with L^p -bounds on curvature, *Commun. Math. Phys.* 83 (1982) 31-42.